



12-1-2023

Sex differences in acute myocardial infarction: clinical characteristics, management practices, and outcomes for patients in a large tertiary hospital from Palestine

Yunis Daralammouri

Department of Medicine, College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine. & Department of Cardiology, An-Najah National University Hospital, Nablus 44839, Palestine

Mohammed Jabri

Department of Medicine, College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine

Osayd Mosleh

Department of Medicine, College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine

Yazan AbdelAziz

Department of Medicine, College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine

Murad Azamtta

*Additional works at: <https://pmpj.najah.edu/journal>
Department of Cardiology, An-Najah National University Hospital, Nablus 44839, Palestine*

Recommended Citation

See next page for additional Authors
Daralammouri, Yunis; Jabri, Mohammed; Mosleh, Osayd; AbdelAziz, Yazan; Azamtta, Murad; Abu Taha, Adham; and Zyoud, Sa'ed (2023) "Sex differences in acute myocardial infarction: clinical characteristics, management practices, and outcomes for patients in a large tertiary hospital from Palestine," *Palestinian Medical and Pharmaceutical Journal*: Vol. 8 : Iss. 3 , Article 15.
Available at: <https://doi.org/10.59049/2790-0231.1257>

This Research article is brought to you for free and open access by Palestinian Medical and Pharmaceutical Journal. It has been accepted for inclusion in Palestinian Medical and Pharmaceutical Journal by an authorized editor of Palestinian Medical and Pharmaceutical Journal. For more information, please contact mqneibi@najah.edu.

Sex differences in acute myocardial infarction: clinical characteristics, management practices, and outcomes for patients in a large tertiary hospital from Palestine

Abstract

Several studies have shown gender variations in acute coronary syndrome's clinical presentation, diagnosis, therapy, and outcomes. Both immediate and long-term outcomes were worse for women with the acute coronary syndrome than men. This study investigates the influence of sex differences on the clinical presentations, treatment approaches, and patient outcomes of acute myocardial infarction at a large tertiary hospital in Palestine. A retrospective cohort study included all patients with acute myocardial infarction who presented to An-Najah National University Hospital from January 2018 to December 2020. Of the 422 patients in this study, 96 (22.7%) were women. Compared to men, women were older ($p < 0.001$) and had significantly higher rates of diabetes mellitus ($p < 0.001$) and hypertension ($p < 0.001$). Additionally, women had greater chances of complaining of atypical cardiac chest pain ($p = 0.012$). Furthermore, non-ST segment elevation myocardial infarction was more prevalent among women ($p = 0.017$). Regarding the hospital course, the median number of stents was statistically significant in men ($p = 0.029$), but women had significantly higher in-hospital mortality ($p = 0.013$) and a higher rate of blood transfusion ($p = 0.026$). Myocardial infarction presents differently in males and women. Women were older, had more comorbid conditions than men, had atypical presentations, and had higher in-hospital mortality rates. As a result, while evaluating and treating patients suspected of having a myocardial infarction, healthcare practitioners must account for these discrepancies to reduce the mortality rate among women.

Keywords

Palestine; Myocardial Infarction; Sex Differences; Presentation; Management; Outcome

Cover Page Footnote

All of the personnel at NNUH's Cardiology Departments merit much praise and gratitude for their enthusiastic participation in this study and active cooperation

Authors

Yunis Daralammouri, Mohammed Jabri, Osayd Mosleh, Yazan AbdelAziz, Murad Azamtta, Adham Abu Taha, and Sa'ed Zyoud

Sex differences in acute myocardial infarction: clinical characteristics, management practices, and outcomes for patients in a large tertiary hospital from Palestine

Yunis Daralammouri^{1, 2, *}, Mohammed Jabri¹, Osayd Mosleh¹, Yazan AbdelAziz¹, Murad Azamtta², Adham Abu Taha^{3, 4} & Sa'ed Zyoud^{5, 6, 7, *}

¹ Department of Medicine, College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine. ² Department of Cardiology, An-Najah National University Hospital, Nablus 44839, Palestine. ³ Department of Pathology, An-Najah National University Hospital, Nablus 44839, Palestine. ⁴ Department of Biomedical Sciences, College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine. ⁵ Department of Clinical and Community Pharmacy, College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine. ⁶ Poison Control and Drug Information Center (PCDIC), College of Medicine and Health Sciences, An-Najah National University, Nablus 44839, Palestine. ⁷ Clinical Research Center, An-Najah National University Hospital, Nablus 44839, Palestine

*Corresponding authors: Yunis71@yahoo.de; saedyoud@yahoo.com

Received: (15/12/2022), Accepted: (12/2/2023)

ABSTRACT

Several studies have shown gender variations in acute coronary syndrome's clinical presentation, diagnosis, therapy, and outcomes. Both immediate and long-term outcomes were worse for women with the acute coronary syndrome than men. This study investigates the influence of sex differences on the clinical presentations, treatment approaches, and patient outcomes of acute myocardial infarction at a large tertiary hospital in Palestine. A retrospective cohort study included all patients with acute myocardial infarction who presented to An-Najah National University Hospital from January 2018 to December 2020. Of the 422 patients in this study, 96 (22.7%) were women. Compared to men, women were older ($p < 0.001$) and had significantly higher rates of diabetes mellitus ($p < 0.001$) and hypertension ($p < 0.001$). Additionally, women had greater chances of complaining of atypical cardiac chest pain ($p = 0.012$). Furthermore, non-ST segment elevation myocardial infarction was more prevalent among women ($p = 0.017$). Regarding the hospital course, the median number of stents was statistically significant in men ($p = 0.029$), but women had significantly higher in-hospital mortality ($p = 0.013$) and a higher rate of blood transfusion ($p = 0.026$). Myocardial infarction presents differently in males and women. Women were older, had more comorbid conditions than men, had atypical presentations, and had higher in-hospital mortality rates. As a result, while evaluating and treating patients suspected of having a myocardial infarction, healthcare practitioners must account for these discrepancies to reduce the mortality rate among women.

Keywords: Palestine; Myocardial Infarction; Sex Differences; Presentation; Management; Outcome.

INTRODUCTION

Cardiovascular disease is the world's leading cause of morbidity and mortality [1, 2]. Acute coronary syndrome (ACS) is a term used to describe a spectrum of clinical presentations that result from acute myocardial ischemia. It includes unstable angina (UA), non-ST-segment elevation myocardial infarction (NSTEMI), and ST-segment elevation myocardial infarction (STEMI). It constitutes a major element of ischemic heart disease [3]. Mortality due to ACS has declined in recent decades due to advances in therapy, lifestyle improvements, and an emphasis on primary prevention, yet high rates remain [4, 5].

Emerging evidence has revealed sex-specific variations in coronary function and anatomy, baseline risk factors, ACS clinical characteristics, pathogenesis, diagnosis, therapeutic efficiency, and outcomes in women [6-11]. Women's underrepresentation exacerbates these significant disparities in cardiovascular clinical studies [12-14].

Women's epicardial coronary arteries are smaller, resulting in increased endothelial shear stress, which might account for gender differences in susceptibility to CAD [15-17]. Low endothelial shear stress has been linked to plaque instability, pathological remodeling, and local lipid accumulation [18].

Furthermore, the phenotype of low vascular resistance and anti-inflammatory activity mediated by estrogen protects against CVD. [19, 20]. Compared to men, women with ACS had even worse short- and long-term results [21-23]. Some predictors include old age at presentation, higher comorbid conditions, and atypical complaints. These atypical complaints might be attributed to delays in diagnosis and treatment, resulting in poor clinical outcomes [8, 9]. However, the mechanisms behind these variations remain largely unexplored [10, 11].

In Palestine, there are insufficient data studying sex differences in acute myocardial infarction. As a result, we opted to investigate the association between sex variations, clinical presentations, treatment processes, and patient outcomes in MI in a large Palestinian tertiary hospital. This study provides new information to help improve morbidity and mortality in our community.

METHODS

Study design, settings, and population

This retrospective study was conducted between January 2018 and December 2020 at An-Najah National University Hospital (NNUH), a large tertiary referral hospital. Its cardiology department is one of the main cardiology centers in the north of West Bank, with three interventional cardiology consultants providing continuous primary percutaneous coronary intervention (PCI) services day and night for patients from all over Palestine. The sample size was all MI patients admitted to An-Najah National University Hospital from January 2018 to December 2020. Age, sex, smoking history, STEMI, NSTEMI, comorbidities (e.g., hypertension, diabetes mellitus, chronic renal disease, cerebrovascular disease, dyslipidemia, and coronary artery disease), prior PCI, and previous coronary artery bypass graft (CABG) were collected and analyzed. The final study group consisted of 422 eligible patients.

Inclusion and exclusion criteria

Inclusion criteria

- Patients were over 18 years old.
- Meet MI criteria.

- Patients have presented or transferred to a large tertiary hospital with the required facilities.

Exclusion criteria

Patients who had MI during hospitalization.

Data collection

We designed a three-section data collection form.

- Section 1 includes demographic characteristics, including age, sex, and smoking status.
- Section 2 includes the type of MI, patient comorbidities, medication, PCI history, and CABG history.
- Section 3 includes clinical presentation, type of intervention made, in-hospital complications related to MI, and in-hospital mortality.

Ethical considerations

The research protocol, which includes the use and access of patient clinical data, was approved by the *Institutional Review Boards (IRB) of An-Najah National University* and the regional health authorities. We confirm that the information gathered was only utilized for clinical research.

Statistical analysis

The data were entered and analyzed using IBM-SPSS version 21, a widely used statistical software package. The data are presented as continuous variables' means and standard deviations (SD), while categorical variables are expressed as frequencies and percentages. The medians were used to represent nonnormally distributed variables (lower-upper quartiles). The variables' normality was then verified using the Kolmogorov–Smirnov test. To examine the significance between categorical variables, exact chi-square or Fisher exact tests were performed as needed. Next, the Kruskal–Wallis or Mann–Whitney tests were used for median differences between categories. A p-value of 0.05 was used to determine the significance threshold.

RESULTS

Four hundred twenty-two patients were enrolled in this study over two years, of whom 96 (22.7%) were women. The baseline features are shown in Table 1, stratified by sex. Compared to men, women were older (mean age 67 vs. 60 years; $p < 0.001$), were significantly less likely to use tobacco (2 vs. 63%; $p < 0.001$), and had significantly more diabetes mellitus (67 vs. 47%; $p < 0.001$) and hypertension (75 vs. 50%; $p < 0.001$). However, there

was no substantial disparity in dyslipidemia between men and women (5 vs. 3%; $p < 0.319$), chronic renal disease (15 vs. 13%; $p < 0.653$), congestive heart failure (8 vs. 10%; $p < 0.547$), or stroke (5 vs. 4%; $p < 0.603$). Regarding the history of CABG and PCI, there were no notable disparities between women and men (6 vs. 9%; $p < 0.283$ and 27 vs. 30%; $p < 0.573$, respectively). More commonly, women who presented to the hospital had atypical chest pain (44 vs. 31%; $p = 0.012$).

Table (1): Baseline characteristics of patients with myocardial infarction according to sex.

| Variable | Male (n=326), n (%) | Female (n=96), n (%) | P value, n (%) |
|-----------------------------|---------------------|----------------------|----------------|
| Age, (mean±SD, years) | 60.02±10.870 | 67.85±10.703 | <0.001 |
| Risk Factors | | | |
| Smoking | 206 (63.2) | 2 (2.1) | <0.001 |
| DM | 154 (47.2) | 65 (67.7) | <0.001 |
| Hypertension | 166 (50.9) | 72 (75) | <0.001 |
| Dyslipidemia | 10 (3.1) | 5 (5.2) | 0.319 |
| History of CABG | 32 (9.8) | 6 (6.3) | 0.283 |
| History of PCI | 98 (30.1) | 26 (27.1) | 0.573 |
| Medical History | | | |
| CKD | 45 (13.8) | 15 (15.6) | 0.653 |
| CHF | 34 (10.4) | 8 (8.3) | 0.547 |
| Stroke | 13 (4) | 5 (5.2) | 0.603 |
| Chest pain character | | | |
| Typical | 225 (69) | 53 (55.2) | 0.012 |
| Atypical | 101 (31) | 43 (44.8) | 0.012 |
| Time , median (Q1-Q3) * | 24 (4-48) | 24 (18-48) | 0.003 |
| Type of MI | | | |
| STEMI | 117 (35.9) | 22 (22.9) | 0.017 |
| NSTEMI | 209 (64.1) | 74 (77.1) | 0.017 |
| Medication history | | | |
| Aspirin | 204 (62.6) | 53 (55.2) | 0.193 |
| Clopidogrel | 47 (14.4) | 11 (11.5) | 0.459 |
| B blocker | 74 (22.7) | 27 (28.1) | 0.273 |
| Statin | 102 (31.3) | 30 (31.3) | 0.994 |
| ACE-I | 97 (29.8) | 27 (28.1) | 0.758 |
| ARB | 41 (12.6) | 12 (12.5) | 0.984 |
| Anticoagulant | 7 (2.1) | 4 (4.2) | 0.281 |
| Laboratory Result | | | |
| Creatinine, median (Q1-Q3) | 0.92 (0.8-1.3) | 0.9 (0.7-1.3375) | 0.181 |
| Troponin, median (Q1-Q3) | 1.7 (0.4-7.2) | 1.285 (0.6-3.335) | 0.412 |
| Echocardiography | | | |
| LVEF, median (Q1-Q3) | 50 (40-55) | 45 (40-55) | 0.582 |

Abbreviations: DM, diabetes mellitus; CABG, coronary artery bypass graft; PCI, percutaneous coronary intervention; CKD, chronic kidney disease; CHF, congestive heart failure; STEMI, ST-segment elevation myocardial infarction; NSTEMI, non-ST segment elevation myocardial infarction; ACE-I, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; LVEF, left ventricular ejection fraction.

*Time from symptoms to the presentation (hours).

Regarding the in-hospital course shown in Table 2, men had a significantly higher median number of stents used than women ($p = 0.029$). In contrast, women were significantly more likely to receive a blood transfusion (4 vs. 0.6%: $p = 0.026$). There were no statistically relevant variations in recurrent angina (0

vs. 0.6%: $p > 0.999$), reinfarction (1 vs. 0.6%: $p = 0.54$), cardiogenic shock (4 vs. 1%: $p = 0.245$), bleeding (1 vs. 0.6%: $p = 0.54$), renal impairment (5 vs. 3%: $p = 0.408$), or cardiac arrest (4 vs. 3%: $p = 0.767$). Women's in-hospital mortality rates were significantly higher than men's (7 vs. 2%: $p = 0.013$).

Table (2): In-hospital course of patients with myocardial infarction according to sex.

| Variable | Male (n=326), n(%) | Female (n=96), n (%) | P value, n (%) |
|---|--------------------|----------------------|----------------|
| Number of coronary vessels, median (Q1-Q3)* | 2 (1-3) | 2 (1-3) | 0.923 |
| Number of stents, median (Q1-Q3) | 1 (0-2) | 1 (0-1) | 0.029 |
| Complications | | | |
| Recurrent angina | 2 (0.6) | 0 (0) | >0.999 |
| Reinfarction | 2 (0.6) | 1 (1) | 0.540 |
| Cardiogenic shock | 6 (1.8) | 4 (4.2) | 0.245 |
| Bleeding | 2 (0.6) | 1 (1) | 0.540 |
| Transfusion | 2 (0.6) | 4 (4.2) | 0.026 |
| Renal impairment | 11 (3.4) | 5 (5.2) | 0.408 |
| Cardiac arrest | 12 (3.7) | 4 (4.2) | 0.767 |
| In-hospital mortality | 7 (2.1) | 7 (7.3) | 0.013 |

*Number of major coronary vessels with significant lesions.

DISCUSSION

Emerging studies over the last decade have revealed sex variations in the clinical manifestations, diagnosis, pathogenesis, and therapy of ACS in women. ACS women were older and had many more comorbidities. They had atypical angina more commonly than men and took far longer to seek medical help. Furthermore, women are more prone to MI-related in-hospital complications and mortality [8, 24-27].

Several agencies have recently advocated for more women to be included in clinical trials investigating gender differences [28-30]. The proportion of women in clinical trials has increased recently, yet sex-specific analyses are still insufficiently undertaken[30]. In our study, the female proportion was 22.7%, less than that of other global studies, reaching almost 38.2% [31].

In this investigation, women with myocardial infarction were older, as in previous research [25, 32, 33]. This might be explained by estrogen's ability to protect against coronary vascular disease, presumably through anti-inflammatory effects and by producing

low vascular resistance in blood vessels [19, 20]. Furthermore, women have increased endothelial shear stress, which protects against CVD by reducing lipid buildup and plaque instability [16-18].

Women reported significantly greater diabetes and hypertension than men in our study, consistent with previous research [8, 34, 35]. This is supported by the fact that there is evidence to suggest that women, on average, may experience more socioeconomic disadvantages and psychosocial stressors than men, which can lead to poorer physical and mental health outcomes and lower quality of life [36, 37]. This is consistent with the observation that women's psychological stress has grown dramatically over the previous two decades as their economic contribution and educational attainment have increased [38].

As in previous research, women are more likely to present with atypical angina and wait longer to seek medical attention [25, 39, 40]. Women are more likely to experience atypical symptoms during a heart attack, such as nausea, vomiting, back or jaw pain, shortness of breath, or extreme fatigue, which may be

mistakenly attributed to other conditions. This can lead to delays in seeking medical attention and receiving appropriate treatment for a heart attack. Additionally, women are more likely to downplay their symptoms or dismiss them as stress- or anxiety-related, which can delay seeking help. Both women and healthcare providers must be aware of these atypical symptoms and take them seriously to ensure timely and appropriate care [41]. As a result, women are at a higher risk of receiving an inaccurate diagnosis and delayed intervention, as proven by multiple studies finding significant system delays in women [42-44]. Furthermore, in some regions of our society, women may have reduced access to healthcare services, which can contribute to delayed presentation. As a result, women presenting with noncardiac symptoms, particularly those with cardiovascular risk factors, require attentive care.

Women are more likely to have NSTEMI as the primary diagnosis and require less angioplasty for therapy with fewer stents, consistent with prior research [42, 43, 45]. Women have a lower proportion of obstructive coronary artery disease and are more likely than men to develop microvascular angina [46, 47]. Others contend that the lack of sex recommendations and the significant proportion of women having nontypical symptoms and comorbid conditions contribute to women's lower likelihood than men of undergoing invasive cardiac intervention [48, 49].

The gender difference in the current study substantially influenced in-hospital complications such as blood transfusion and in-hospital mortality, which are greater in women. This finding is consistent with earlier research [25, 33, 43, 50-52]. Higher blood transfusion rates in women have been linked to advanced age, comorbidities, body weight, and homeostasis variations due to the menstrual cycle, hormonal contraceptives or hormonal replacement therapy (HRT), menopause, and pregnancy [53, 54]. Furthermore, estrogen's direct inhibitory action on platelet aggregation may contribute to women's higher bleeding risk [46, 47, 55].

Multiple factors might contribute to in-hospital mortality. For instance, women with ACS are often older and have more concomitant diseases than males. Furthermore, women are less likely than males to have chest pain,

diagnostic electrocardiograph (ECG) abnormalities, or elevated troponin levels on arrival, resulting in an incorrect diagnosis and treatment delay [56-58]. Finally, women have more mental stress after ACS than men, which leads to worse results [47, 59-61].

This emphasizes the need for social health practitioners to perform more comprehensive assessments in hospitals and after discharge to enhance the psychosocial domain [47].

Study limitations

This study was carried out at a single center (NNUH), and the study population was comparatively small to examine all variables. Our data are based on an observational retrospective registry and did not include patients with acute myocardial infarction who had not been admitted to the hospital. Moreover, the female proportion in our study was less than that in other global studies, which could affect the results of our study. Other demographic characteristics, such as education level, geographical area, and economic status, were not considered. However, well-designed registry data will provide valid results.

CONCLUSIONS

Myocardial infarction manifests differently in men and women. Women had more comorbidities than males with an initial diagnosis of myocardial infarction. Women were often older than males and had higher in-hospital mortality rates. This increase in female mortality rates may be due to women having more atypical symptoms, resulting in misdiagnosis and undertreatment. We advocate that healthcare practitioners consider these disparities when diagnosing and treating patients suspected of having a myocardial infarction to reduce the fatality rate among women. We also suggest that future studies on myocardial infarction be performed in Palestine, given that there are preliminary studies on this life-threatening condition.

Ethics approval and consent to participate

The administrator of An-Najah National University Hospital and the institution's institutional review board (IRB) approved this study. The study methodologies were all carried out under the required laws and guidelines. Since we used retrospective data, An-

Najah National University's IRB omitted the requirement for informed consent.

Consent to publish

Not applicable

Availability of data and materials

Because of the ethical approval requirements regarding patient data and confidentiality, the datasets created and/or analyzed during the current research are not publicly available but accessible from the corresponding author upon justifiable inquiry. The abstract was released as part of institutional repositories' self-archiving initiatives (i.e., university repository).

Competing interests

The authors state that they have no conflicts of interest.

FUNDING

None

Authors' contributions

MSJ, OSM, and YNA performed data collection, data analysis, and manuscript drafting after reviewing the literature. MA and YD participated in the study's design, assessed patients for eligibility, reviewed the literature, and provided an important manuscript revision. AAT drafted the manuscript, was responsible for the integrity of the data, and conducted a critical assessment of the research to improve its intellectual quality. The research was developed and designed by SHZ, who also oversaw, coordinated, and analyzed the data, provided critical feedback on interpreting the findings, and helped with the final manuscript writing. The final manuscript was then read by all authors and approved. This paper was included in a graduation project for the Doctor of Medicine program at An-Najah National University.

ACKNOWLEDGMENT

All of the personnel at NNUH's Cardiology Departments merit much praise and gratitude for their enthusiastic participation in this study and active cooperation.

REFERENCES

- 1] Nowbar AN, Gitto M, Howard JP, Francis DP, Al-Lamee R. Mortality From Ischemic Heart Disease. *Circ Cardiovasc Qual Outcomes*. 2019;12(6):e005375.
- 2] Vedanthan R, Seligman B, Fuster V. Global perspective on acute coronary syndrome: a burden on the young and poor. *Circ Res*. 2014;114(12):1959-75.
- 3] An L, Li W, Shi H, Zhou X, Liu X, Wang H, Liu J, Fan S. Gender difference of symptoms of acute coronary syndrome among Chinese patients: a cross-sectional study. *Eur J Cardiovasc Nurs*. 2019;18(3):179-84.
- 4] Roth GA, Johnson C, Abajobir A, Abd-Allah F, Abera SF, Abyu G, Ahmed M, Aksut B, Alam T, Alam K, Alla F, Alvis-Guzman N, Amrock S, Ansari H, Ärnlöv J, Asayesh H, Atey TM, Avila-Burgos L, Awasthi A, Banerjee A, Barac A, Barnighausen T, Barregard L, Bedi N, Belay Ketema E, Bennett D, Berhe G, Bhutta Z, Bitew S, Carapetis J, Carrero JJ, Malta DC, Castañeda-Orjuela CA, Castillo-Rivas J, Catalá-López F, Choi JY, Christensen H, Cirillo M, Cooper L, Jr., Criqui M, Cundiff D, Damasceno A, Dandona L, Dandona R, Davletov K, Dharmaratne S, Dorairaj P, Dubey M, Ehrenkranz R, El Sayed Zaki M, Faraon EJA, Esteghamati A, Farid T, Farvid M, Feigin V, Ding EL, Fowkes G, Gebrehiwot T, Gillum R, Gold A, Gona P, Gupta R, Habtewold TD, Hafezi-Nejad N, Hailu T, Hailu GB, Hankey G, Hassen HY, Abate KH, Havmoeller R, Hay SI, Horino M, Hotez PJ, Jacobsen K, James S, Javanbakht M, Jeemon P, John D, Jonas J, Kalkonde Y, Karimkhani C, Kasaeian A, Khader Y, Khan A, Khang YH, Khera S, Khoja AT, Khubchandani J, Kim D, Kolte D, Kosen S, Krohn KJ, Kumar GA, Kwan GF, Lal DK, Larsson A, Linn S, Lopez A, Lotufo PA, El Razek HMA, Malekzadeh R, Mazidi M, Meier T, Meles KG, Mensah G, Meretoja A, Mezgebe H, Miller T, Mirrahimov E, Mohammed S, Moran AE, Musa KI, Narula J, Neal B, Ngalesoni F, Nguyen G, Obermeyer CM, Owolabi M, Patton G, Pedro J, Qato D, Qorbani M, Rahimi K, Rai RK, Rawaf S, Ribeiro A, Safiri S, Salomon JA, Santos I, Santric Milicevic M, Sartorius B, Schutte A, Sepanlou S, Shaikh MA, Shin MJ, Shishehbor M, Shore H, Silva DAS, Sobngwi E, Stranges S, Swaminathan S, Tabarés-

- Seisedos R, Tadele Atnafu N, Tesfay F, Thakur JS, Thrift A, Topor-Madry R, Truelsen T, Tyrovolas S, Ukwaja KN, Uthman O, Vasankari T, Vlassov V, Vollset SE, Wakayo T, Watkins D, Weintraub R, Werdecker A, Westerman R, Wiysonge CS, Wolfe C, Workicho A, Xu G, Yano Y, Yip P, Yonemoto N, Younis M, Yu C, Vos T, Naghavi M, Murray C. Global, Regional, and National Burden of Cardiovascular Diseases for 10 Causes, 1990 to 2015. *J Am Coll Cardiol*. 2017;70(1):1-25.
- 5] Koopman C, Vaartjes I, Heintjes EM, Spiering W, van Dis I, Herings RM, Bots ML. Persisting gender differences and attenuating age differences in cardiovascular drug use for prevention and treatment of coronary heart disease, 1998-2010. *Eur Heart J*. 2013;34(41):3198-205.
- 6] Hambraeus K, Tydén P, Lindahl B. Time trends and gender differences in prevention guideline adherence and outcome after myocardial infarction: Data from the SWEDEHEART registry. *Eur J Prev Cardiol*. 2016;23(4):340-8.
- 7] Sabbag A, Matetzky S, Porter A, Iakobishvili Z, Moriel M, Zwas D, Fefer P, Asher E, Beigel R, Gottlieb S, Goldenberg I, Segev A. Sex Differences in the Management and 5-Year Outcome of Young Patients (<55 Years) with Acute Coronary Syndromes. *Am J Med*. 2017;130(11):1324.e15-e22.
- 8] Shehab A, Al-Dabbagh B, AlHabib KF, Alsheikh-Ali AA, Almahmeed W, Sulaiman K, Al-Motarreb A, Nagelkerke N, Al Suwaidi J, Hersi A, Al Faleh H, Asaad N, Al Saif S, Amin H. Gender disparities in the presentation, management and outcomes of acute coronary syndrome patients: data from the 2nd Gulf Registry of Acute Coronary Events (Gulf RACE-2). *PLoS One*. 2013;8(2):e55508.
- 9] Dey S, Flather MD, Devlin G, Brieger D, Gurfinkel EP, Steg PG, Fitzgerald G, Jackson EA, Eagle KA. Sex-related differences in the presentation, treatment and outcomes among patients with acute coronary syndromes: the Global Registry of Acute Coronary Events. *Heart*. 2009;95(1):20-6.
- 10] Vaccarino V. Myocardial Infarction in Young Women. *Circulation*. 2019;139(8):1057-9.
- 11] Pagidipati NJ, Peterson ED. Acute coronary syndromes in women and men. *Nat Rev Cardiol*. 2016;13(8):471-80.
- 12] Garcia M, Mulvagh SL, Merz CN, Buring JE, Manson JE. Cardiovascular Disease in Women: Clinical Perspectives. *Circ Res*. 2016;118(8):1273-93.
- 13] Scott PE, Unger EF, Jenkins MR, Southworth MR, McDowell TY, Geller RJ, Elahi M, Temple RJ, Woodcock J. Participation of Women in Clinical Trials Supporting FDA Approval of Cardiovascular Drugs. *J Am Coll Cardiol*. 2018;71(18):1960-9.
- 14] Ghare MI, Chandrasekhar J, Mehran R, Ng V, Grines C, Lansky A. Sex Disparities in Cardiovascular Device Evaluations: Strategies for Recruitment and Retention of Female Patients in Clinical Device Trials. *JACC Cardiovasc Interv*. 2019;12(3):301-8.
- 15] Hiteshi AK, Li D, Gao Y, Chen A, Flores F, Mao SS, Budoff MJ. Gender differences in coronary artery diameter are not related to body habitus or left ventricular mass. *Clin Cardiol*. 2014;37(10):605-9.
- 16] Patel MB, Bui LP, Kirkeeide RL, Gould KL. Imaging Microvascular Dysfunction and Mechanisms for Female-Male Differences in CAD. *JACC Cardiovasc Imaging*. 2016;9(4):465-82.
- 17] Kerkhof PL, Miller VM. Sex-Specific analysis of cardiovascular function: Springer; 2018.
- 18] Koskinas KC, Sukhova GK, Baker AB, Papafaklis MI, Chatzizisis YS, Coskun AU, Quillard T, Jonas M, Maynard C, Antoniadis AP, Shi GP, Libby P, Edelman ER, Feldman CL, Stone PH. Thin-capped atheromata with reduced collagen content in pigs develop in coronary arterial regions exposed to persistently low endothelial shear stress. *Arterioscler Thromb Vasc Biol*. 2013;33(7):1494-504.

- 19] Deroo BJ, Korach KS. Estrogen receptors and human disease. *J Clin Invest.* 2006;116(3):561-70.
- 20] Miller VM, Duckles SP. Vascular actions of estrogens: functional implications. *Pharmacol Rev.* 2008;60(2):210-41.
- 21] Shehab A, AlHabib KF, Bhagavathula AS, Hersi A, Alfaleh H, Alshamiri MQ, Ullah A, Sulaiman K, Almahmeed W, Al Suwaidi J, Alsheikh-Ali AA, Amin H, Al Jarallah M, Salam AM. Clinical Presentation, Quality of Care, Risk Factors and Outcomes in Women with Acute ST-Elevation Myocardial Infarction (STEMI): An Observational Report from Six Middle Eastern Countries. *Curr Vasc Pharmacol.* 2019;17(4):388-95.
- 22] D'Onofrio G, Safdar B, Lichtman JH, Strait KM, Dreyer RP, Geda M, Spertus JA, Krumholz HM. Sex differences in reperfusion in young patients with ST-segment-elevation myocardial infarction: results from the VIRGO study. *Circulation.* 2015;131(15):1324-32.
- 23] Piackova E, Jäger B, Farhan S, Christ G, Schreiber W, Weidinger F, Stefanelli T, Delle-Karth G, Kaff A, Maurer G, Huber K. Gender differences in short- and long-term mortality in the Vienna STEMI registry. *Int J Cardiol.* 2017;244:303-8.
- 24] El-Menyar A, Zubaid M, Sulaiman K, AlMahmeed W, Singh R, Alsheikh-Ali AA, Al Suwaidi J. Atypical presentation of acute coronary syndrome: a significant independent predictor of in-hospital mortality. *J Cardiol.* 2011;57(2):165-71.
- 25] Khesroh AA, Al-Roumi F, Al-Zakwani I, Attur S, Rashed W, Zubaid M. Gender Differences among Patients with Acute Coronary Syndrome in the Middle East. *Heart Views.* 2017;18(3):77-82.
- 26] El-Menyar A, Zubaid M, Rashed W, Almahmeed W, Al-Lawati J, Sulaiman K, Al-Motarreb A, Amin H, R S, Al Suwaidi J. Comparison of men and women with acute coronary syndrome in six Middle Eastern countries. *Am J Cardiol.* 2009;104(8):1018-22.
- 27] Steg PG, Goldberg RJ, Gore JM, Fox KA, Eagle KA, Flather MD, Sadiq I, Kasper R, Rushton-Mellor SK, Anderson FA. Baseline characteristics, management practices, and in-hospital outcomes of patients hospitalized with acute coronary syndromes in the Global Registry of Acute Coronary Events (GRACE). *Am J Cardiol.* 2002;90(4):358-63.
- 28] Schiebinger L, Leopold SS, Miller VM. Editorial policies for sex and gender analysis. *Lancet.* 2016;388(10062):2841-2.
- 29] Avery E, Clark J. Sex-related reporting in randomised controlled trials in medical journals. *Lancet.* 2016;388(10062):2839-40.
- 30] Liu KA, Mager NA. Women's involvement in clinical trials: historical perspective and future implications. *Pharm Pract (Granada).* 2016;14(1):708.
- 31] Jin X, Chandramouli C, Allocco B, Gong E, Lam CSP, Yan LL. Women's Participation in Cardiovascular Clinical Trials From 2010 to 2017. *Circulation.* 2020;141(7):540-8.
- 32] Shehab A, Bhagavathula AS, Alhabib KF, Ullah A, Suwaidi JA, Almahmeed W, AlFaleh H, Zubaid M. Age-Related Sex Differences in Clinical Presentation, Management, and Outcomes in ST-Segment-Elevation Myocardial Infarction: Pooled Analysis of 15 532 Patients From 7 Arabian Gulf Registries. *J Am Heart Assoc.* 2020;9(4):e013880.
- 33] Shehab A, Yasin J, Hashim MJ, Al-Dabbagh B, Mahmeed WA, Bustani N, Agrawal A, Yusufali A, Wassef A, Alnaeemi A. Gender differences in acute coronary syndrome in Arab Emirati women--implications for clinical management. *Angiology.* 2013;64(1):9-14.
- 34] Elsaesser A, Hamm CW. Acute coronary syndrome: the risk of being female. *Circulation.* 2004;109(5):565-7.
- 35] Maas AH, Appelman YE. Gender differences in coronary heart disease. *Neth Heart J.* 2010;18(12):598-602.
- 36] Buchholz EM, Strait KM, Dreyer RP, Lindau ST, D'Onofrio G, Geda M, Spatz ES, Beltrame JF, Lichtman JH, Lorenze NP, Bueno H, Krumholz HM. Editor's

- Choice-Sex differences in young patients with acute myocardial infarction: A VIRGO study analysis. *Eur Heart J Acute Cardiovasc Care*. 2017;6(7):610-22.
- 37] Allabadi H, Probst-Hensch N, Alkaiyat A, Haj-Yahia S, Schindler C, Kwiatkowski M, Zemp E. Mediators of gender effects on depression among cardiovascular disease patients in Palestine. *BMC Psychiatry*. 2019;19(1):284.
- 38] Schwab K, Samans R, Zahidi S, et al. The global gender gap report 2016: World Economic Forum; 2016.
- 39] Mehta LS, Beckie TM, DeVon HA, Grines CL, Krumholz HM, Johnson MN, Lindley KJ, Vaccarino V, Wang TY, Watson KE, Wenger NK. Acute Myocardial Infarction in Women: A Scientific Statement From the American Heart Association. *Circulation*. 2016;133(9):916-47.
- 40] Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, Bax JJ, Borger MA, Brotons C, Chew DP, Gencer B, Hasenfuss G, Kjeldsen K, Lancellotti P, Landmesser U, Mehilli J, Mukherjee D, Storey RF, Windecker S. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2016;37(3):267-315.
- 41] Lichtman JH, Leifheit EC, Safdar B, Bao H, Krumholz HM, Lorenze NP, Daneshvar M, Spertus JA, D'Onofrio G. Sex Differences in the Presentation and Perception of Symptoms Among Young Patients With Myocardial Infarction: Evidence from the VIRGO Study (Variation in Recovery: Role of Gender on Outcomes of Young AMI Patients). *Circulation*. 2018;137(8):781-90.
- 42] Gebhard CE, Gebhard C, Maafi F, Bertrand MJ, Stähli BE, Maredziak M, Bengs S, Haider A, Zhang ZW, Smith DC, Ly HQ. Impact of summer season on pre-hospital time delays in women and men undergoing primary percutaneous coronary intervention. *Sci Total Environ*. 2019;656:322-30.
- 43] Meyer MR, Bernheim AM, Kurz DJ, O'Sullivan CJ, Tüller D, Zbinden R, Rosemann T, Eberli FR. Gender differences in patient and system delay for primary percutaneous coronary intervention: current trends in a Swiss ST-segment elevation myocardial infarction population. *Eur Heart J Acute Cardiovasc Care*. 2019;8(3):283-90.
- 44] Roswell RO, Kunkes J, Chen AY, Chiswell K, Iqbal S, Roe MT, Bangalore S. Impact of Sex and Contact-to-Device Time on Clinical Outcomes in Acute ST-Segment Elevation Myocardial Infarction-Findings From the National Cardiovascular Data Registry. *J Am Heart Assoc*. 2017;6(1).
- 45] Zheng H, Foo LL, Tan HC, Richards AM, Chan SP, Lee CH, Low AFH, Hausenloy DJ, Tan JWC, Sahlen AO, Ho HH, Chai SC, Tong KL, Tan DSY, Yeo KK, Chua TSJ, Lam CSP, Chan MY. Sex Differences in 1-Year Rehospitalization for Heart Failure and Myocardial Infarction After Primary Percutaneous Coronary Intervention. *Am J Cardiol*. 2019;123(12):1935-40.
- 46] Jespersen L, Hvelplund A, Abildstrøm SZ, Pedersen F, Galatius S, Madsen JK, Jørgensen E, Kelbæk H, Prescott E. Stable angina pectoris with no obstructive coronary artery disease is associated with increased risks of major adverse cardiovascular events. *Eur Heart J*. 2012;33(6):734-44.
- 47] Haider A, Bengs S, Luu J, Osto E, Siller-Matula JM, Muka T, Gebhard C. Sex and gender in cardiovascular medicine: presentation and outcomes of acute coronary syndrome. *Eur Heart J*. 2020;41(13):1328-36.
- 48] Worrall-Carter L, McEvedy S, Wilson A, Rahman MA. Gender Differences in Presentation, Coronary Intervention, and Outcomes of 28,985 Acute Coronary Syndrome Patients in Victoria, Australia. *Womens Health Issues*. 2016;26(1):14-20.

- 49] Bugiardini R, Yan AT, Yan RT, Fitchett D, Langer A, Manfrini O, Goodman SG. Factors influencing underutilization of evidence-based therapies in women. *Eur Heart J*. 2011;32(11):1337-44.
- 50] Townsend N, Wilson L, Bhatnagar P, Wickramasinghe K, Rayner M, Nichols M. Cardiovascular disease in Europe: epidemiological update 2016. *Eur Heart J*. 2016;37(42):3232-45.
- 51] Berg J, Björck L, Nielsen S, Lappas G, Rosengren A. Sex differences in survival after myocardial infarction in Sweden, 1987-2010. *Heart*. 2017;103(20):1625-30.
- 52] Cenko E, Yoon J, Kedev S, Stankovic G, Vasiljevic Z, Krljanac G, Kalpak O, Ricci B, Milicic D, Manfrini O, van der Schaar M, Badimon L, Bugiardini R. Sex Differences in Outcomes After STEMI: Effect Modification by Treatment Strategy and Age. *JAMA Intern Med*. 2018;178(5):632-9.
- 53] Rubini Gimenez M, Zeymer U, Desch S, de Waha-Thiele S, Ouarrak T, Poess J, Meyer-Saraei R, Schneider S, Fuernau G, Stepinska J, Huber K, Windecker S, Montalescot G, Savonitto S, Jeger RV, Thiele H. Sex-Specific Management in Patients With Acute Myocardial Infarction and Cardiogenic Shock: A Substudy of the CULPRIT-SHOCK Trial. *Circ Cardiovasc Interv*. 2020;13(3):e008537.
- 54] Fu WX, Zhou TN, Wang XZ, Zhang L, Jing QM, Han YL. Sex-Related Differences in Short- and Long-Term Outcome among Young and Middle-Aged Patients for ST-Segment Elevation Myocardial Infarction Underwent Percutaneous Coronary Intervention. *Chin Med J (Engl)*. 2018;131(12):1420-9.
- 55] Renda G, Patti G, Lang IM, Siller-Matula JM, Hylek EM, Ambrosio G, Haas S, De Caterina R. Thrombotic and hemorrhagic burden in women: Gender-related issues in the response to antithrombotic therapies. *Int J Cardiol*. 2019;286:198-207.
- 56] Li S, Fonarow GC, Mukamal KJ, Liang L, Schulte PJ, Smith EE, DeVore A, Hernandez AF, Peterson ED, Bhatt DL. Sex and Race/Ethnicity-Related Disparities in Care and Outcomes After Hospitalization for Coronary Artery Disease Among Older Adults. *Circ Cardiovasc Qual Outcomes*. 2016;9(2 Suppl 1):S36-44.
- 57] Cenko E, van der Schaar M, Yoon J, Kedev S, Valvukis M, Vasiljevic Z, Ašanin M, Miličić D, Manfrini O, Badimon L, Bugiardini R. Sex-Specific Treatment Effects After Primary Percutaneous Intervention: A Study on Coronary Blood Flow and Delay to Hospital Presentation. *J Am Heart Assoc*. 2019;8(4):e011190.
- 58] Hao Y, Liu J, Liu J, Yang N, Smith SC, Jr., Huo Y, Fonarow GC, Ge J, Taubert KA, Morgan L, Zhou M, Xing Y, Ma CS, Han Y, Zhao D. Sex Differences in In-Hospital Management and Outcomes of Patients With Acute Coronary Syndrome. *Circulation*. 2019;139(15):1776-85.
- 59] Xu X, Bao H, Strait K, Spertus JA, Lichtman JH, D'Onofrio G, Spatz E, Bucholz EM, Geda M, Lorenze NP, Bueno H, Beltrame JF, Krumholz HM. Sex differences in perceived stress and early recovery in young and middle-aged patients with acute myocardial infarction. *Circulation*. 2015;131(7):614-23.
- 60] Rutledge T, Kenkre TS, Thompson DV, Bittner VA, Whittaker K, Eastwood JA, Eteiba W, Cornell CE, Krantz DS, Pepine CJ, Johnson BD, Handberg EM, Bairey Merz CN. Psychosocial predictors of long-term mortality among women with suspected myocardial ischemia: the NHLBI-sponsored Women's Ischemia Syndrome Evaluation. *J Behav Med*. 2016;39(4):687-93.
- 61] Vaccarino V, Sullivan S, Hammadah M, Wilmot K, Al Mheid I, Ramadan R, Elon L, Pimple PM, Garcia EV, Nye J, Shah AJ, Alkhoder A, Levantsevych O, Gay H, Obideen M, Huang M, Lewis TT, Bremner JD, Quyyumi AA, Raggi P. Mental Stress-Induced-Myocardial Ischemia in Young Patients With Recent Myocardial Infarction: Sex Differences and Mechanisms. *Circulation*. 2018;137(8):794-805.